

Correction

Open Access

Correction: five-year predictors of physical activity decline among adults in low-income communities: a prospective study

Deborah R Weiss^{1,3}, Jennifer L O'Loughlin², Robert W Platt^{3,4,5} and Gilles Paradis^{*3,6}

Address: ¹Centre for Clinical Epidemiology and Community Studies, Jewish General Hospital, Montréal, Québec, Canada, ²Department of Social and Preventive Medicine, Centre de recherche CHUM, Université de Montréal, Montréal, Québec, Canada, ³Department of Epidemiology, Biostatistics and Occupational Health, Faculty of Medicine, McGill University, Montréal, Québec, Canada, ⁴Montreal Children's Hospital Research Institute, Montréal, Québec, Canada, ⁵Department of Pediatrics, McGill University, Montréal, Québec, Canada and ⁶Institut national de santé publique du Québec, Montreal, Québec, Canada

Email: Deborah R Weiss - deborah.weiss@mail.mcgill.ca; Jennifer L O'Loughlin - jennifer.oloughlin@umontreal.ca; Robert W Platt - robert.platt@mcgill.ca; Gilles Paradis* - gilles.paradis@mcgill.ca

* Corresponding author

Published: 5 June 2007

Received: 15 May 2007

International Journal of Behavioral Nutrition and Physical Activity 2007, **4**:23 doi:10.1186/1479-5868-4-23

Accepted: 5 June 2007

This article is available from: <http://www.ijbnpa.org/content/4/1/23>

© 2007 Weiss et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

After publication it was brought to our attention that the information for one of the variables in Table 1 was incorrect (Weiss, O'Loughlin et al. *International Journal of Behavioral Nutrition and Physical Activity* 2007, **4**:2). The variable in question is "Use of a neighborhood facility for activity". In the first column, the first row should read "yes", and the second row, "no". In the second column, the first row should read 25.8 (41) and the second row, 41.3 (152).

Table 1: Unadjusted and adjusted Odds Ratios for potential predictors of becoming inactive [1].

| Potential predictor | Participants who became inactive | Unadjusted OR (95% CI) | Adjusted OR (95% CI) |
|--|----------------------------------|------------------------|----------------------|
| Sex, % (n) | | | |
| Male | 31.1 (77) | 1.00 (Ref.) | 1.00 (Ref.) |
| Female | 41.6 (116) | 1.58 (1.10, 2.27) | 1.63 (1.09, 2.43) |
| Age (years), mean (SD) | 39.5 (12.2) | 1.03 (1.01, 1.05) | 1.02 (1.01, 1.04) |
| Self-rated health, % (n) | | | |
| Excellent | 29.8 (50) | 1.70 (1.35, 2.16) | 1.39 (1.05, 1.84) |
| Good | 32.6 (84) | | |
| Average | 57.8 (48) | | |
| Poor/Very poor | 64.7 (11) | | |
| BMI ^a , % (n) | | | |
| <25 | 32.1 (117) | 1.00 (Ref.) | 1.00 (Ref.) |
| ≥25 | 46.6 (76) | 1.84 (1.26, 2.69) | 1.57 (1.03, 2.40) |
| Smoking status, % (n) | | | |
| Current smoker | 37.5 (77) | 1.00 (Ref.) | 1.00 (Ref.) |
| Past/Never smoker | 36.2 (121) | 1.06 (0.73, 1.53) | 0.98 (0.78, 1.23) |
| Income, % (n) | | | |
| <20,000 | 39.5 (68) | 0.82 (0.65, 1.03) | 0.95 (0.73, 1.24) |
| 20,000–40,000 | 35.8 (62) | | |
| 40,000+ | 30.2 (42) | | |
| Self-efficacy score, mean (SD) | 2.2 (0.6) | 1.61 (1.20, 2.26) | 1.46 (1.00, 2.14) |
| Use of a neighborhood facility for activity, % (n) | | | |
| Yes | 25.8 (41) | 1.00 (Ref.) | 1.00 (Ref.) |
| No | 41.3 (152) | 2.03 (1.34, 3.06) | 1.61 (1.02, 2.55) |
| Education, % (n) | | | |
| Elementary/some secondary | 52.6 (50) | 0.74 (0.63, 0.86) | N/A* |
| Completed secondary/some college | 40.2 (47) | | |
| Completed college | 32.5 (27) | | |
| Some post-secondary | 29.8 (67) | | |
| Receives encouragement for activity, % (n) | | | |
| No | 35.2 (113) | 1.00 (Ref.) | N/A* |
| Yes | 38.5 (79) | 0.87 (0.60, 1.25) | |

^aBody Mass Index

* Not included in final model

References

1. Weiss DR, O'Loughlin JL, Platt RW, Paradis G: **Five-year predictors of physical activity decline among adults in low-income communities: a prospective study.** *Int J Behav Nutr Phys Act* 4:2. 2007 Jan 18

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

